

Association between level of anxiety and degree of psychosomatic features in medical students at a private university in Northern Peru

Karolay Tacto-Solis,¹ Elizabeth Carolina Muñoz Arteaga,² Jessenia Fiestas-Cordova,² Christian Alberto Rodríguez-Saldaña¹

¹ Facultad de Medicina Humana, Universidad Privada Antenor Orrego, Piura, Perú.

² Facultad de Medicina Humana, Universidad Cesar Vallejo, Piura, Perú.

Correspondence:

Karolay Yanais Tacto-Solis
Facultad de Medicina Humana, Universidad Privada Antenor Orrego, Piura, Perú.
Calle El Rosario 657,
Urbanización Santa Rosa – Sullana.
Phone: +51 918 520 - 819
Email: tactosolik@gmail.com

Received: 10 June 2022

Accepted: 11 October 2022

Citation:

Tacto-Solis, K., Muñoz Arteaga, E. C., Fiestas-Cordova, J., & Rodríguez-Saldaña, C. A. (2023). Association between level of anxiety and degree of psychosomatic features in medical students at a private university in Northern Peru. *Salud Mental*, 46(2), 55-59.

DOI: 10.17711/SM.0185-3325.2023.008



ABSTRACT

Introduction. University students experience multiple academic and social activities that cause enormous anxiety and stress, affecting them psychologically and physically. In this case, the psychosomatic symptoms, resulting from stress and anxiety, are part of a physical ailment and their emergence can be linked to the setting and time when they occur. **Objective.** To determine the association between anxiety levels and the degree of psychosomatic features (PF). **Method.** A study with a cross-sectional analytical design was developed, which included a total of 352 medical students from the city of Piura, Peru, to whom the PHQ-15 test was administered through Google Forms. Those with a previous psychiatric diagnosis were excluded. **Results.** It was found that PF are influenced by being female (RPa = 1.45, CI 95% = [1.23, 1.71], $p \leq .001$), and having mild (RPa = 1.11, CI 95% = [1.20, 1.63], $p \leq .001$) and moderate anxiety levels (RPa = 1.7, CI 95% = [1.24, 2.34], $p = .001$). **Discussion and conclusion.** The presence of a mental condition is necessary for the presence of PF. These stressors create selective alterations of large-scale brain networks involved in the cognitive control, regulation and processing of emotions, stress, and somatic-visceral perception. The study concludes that psychosomatic features are found in approximately 75% of medical students and anxiety in approximately 15%. Mild and moderate anxiety is an influential factor in psychosomatic features, as is being female.

Keywords: Psychophysiological disorders, anxiety, medical students, PHQ-15.

RESUMEN

Introducción. Los estudiantes universitarios viven un escenario de acumulación de actividades académicas y sociales que generan gran ansiedad y estrés, afectándolos psicológica y físicamente. En este caso, los síntomas psicósomáticos, producto del estrés y la ansiedad, forman parte de una dolencia física y pueden vincular su aparición con el escenario y el tiempo que los presenta. **Objetivo.** Determinar la asociación entre el nivel de ansiedad y el grado de manifestación psicósomática. **Método.** Se desarrolló un estudio con diseño analítico transversal que incluyó un total de 352 estudiantes de medicina de la ciudad de Piura, Perú, a quienes se les aplicó mediante Google forms el test de PHQ-15. Se excluyeron a aquellos con diagnóstico psiquiátrico previo. **Resultados.** Se encontró que las MS se ven influenciadas por el sexo femenino (RPa = 1.45, IC 95% = [1.23, 1.71], $p \leq .001$), nivel de ansiedad leve (RPa = 1.11, IC 95% = [1.20, 1.63], $p \leq .001$) y nivel de ansiedad moderada (RPa = 1.7 IC 95% = [1.24, 2.34]; $p = .001$). **Discusión y conclusión.** Es necesario la presencia de una afección mental para la presencia de la MS. Estos factores estresantes generan alteraciones selectivas de redes cerebrales a gran escala implicadas en el control cognitivo, regulación y procesamiento de emociones, estrés y percepción somático-visceral. Se concluye que las manifestaciones psicósomáticas se encuentran en 75% y la ansiedad en 15% aproximadamente, en estudiantes de medicina. La ansiedad leve y moderada es un factor influyente en las manifestaciones psicósomáticas, así como el sexo femenino.

Palabras clave: Trastornos psicofisiológicos, ansiedad, estudiantes de medicina, PHQ-15.

INTRODUCTION

The current global prevalence estimate for anxiety is 26.9% (Nochaiwong et al., 2021). Anxiety is a response of the autonomic nervous system consisting of defensive reactions to danger, which perform a protective function and are considered to be the basis of learning and motivation. If it happens repeatedly, prolonged manner, it disrupts physiological and psychosocial functioning (Galiano-Ramírez, Castellanos-Luna, & Moreno-Mora, 2016). Compounded by the context of the COVID-19 pandemic and the new normality, it leads to psychosomatic disorders (Bartmann et al., 2021; Tollos, Theodorakopoulou, & Christodoulou, 2021).

A psychosomatic disorder is a condition in which psychological stresses negatively affect a person's physiological (somatic) functioning. Psychosomatic symptoms therefore arise as a physiological concomitant of an emotional state (Sajid, Ahmad, & Khalid, 2015). In this case, psychosomatic symptoms, resulting from stress and anxiety, are part of a physical ailment and their emergence can be linked to the setting and time in which they occur (Trebin, 2020).

University students often report anxiety and stress together with psychosomatic symptoms. It has been documented that they may have similar levels to those of other populations, such as the chronically ill. University students are considered to be a particularly vulnerable population, which is exposed to stressors and their effects (Maria-Jereward & Bhuvanewari, 2019).

Medical students are not exempt from this, undergoing adaptive physiological, neuroendocrine, immunological, emotional, and behavioral processes during their university education (Atta & Almilaibary, 2022; Salam et al., 2015). They experience multiple academic and social activities that create enormous anxiety and stress, affecting optimal learning and academic performance (Alzahrani, Alghamdi, Alqarni, Alshareef, & Alzahrani, 2019; Cabanach, Souto-Gestal, & Fernández Cervantes, 2017).

The academic factors associated with the development of anxiety and depression are grade retention, age, and sex (de La Rosa-Rojas et al., 2015; Quito-Calle, Tamayo-Piedra, Buñay-Barahona, & Neira-Cardenas, 2017). Anxiety and depression increase in intensity according to the year of study, with higher levels of anxiety occurring during clinical science semesters (Anton et al., 2021).

It is important to evaluate mental health in medical students, specifically during the pandemic. As a result of the factors mentioned earlier, the aim was to determine the association between anxiety levels and the degree of psychosomatic features in medical students.

METHOD

Study design

A cross-sectional, analytical study was undertaken with convenience sampling.

Subjects/sample description

The study was conducted in the city of Piura in Northern Peru. Three hundred and fifty-two medical students from Antenor Orrego Private University (UPAO), César Vallejo University (UCV) and the National University of Piura (UNP) were enrolled in the study. The UPAO and UCV are private universities, whereas the UNP is a state university. University students enrolled in the 2022-I semester were included. Study participants were youths aged seventeen and older. Those with psychiatric diagnoses prior to their admission to university were excluded.

Measurements

A survey was designed in Google Forms with the instruments used, comprising three sections. The first records general characteristics (age, sex, semester, and grade retention); the second psychosomatic features and the third the degree of anxiety. To assess psychosomatic features, the 15-item PHQ-15 (Patient Health Questionnaire Physical Symptoms) was used, with scores ranging from 0 to 30 and scores ≥ 5 , ≥ 10 , and ≥ 15 representing mild, moderate, and severe levels of somatization (Kroenke, Spitzer, & Williams, 2002). This scale was previously used in a national study on adolescents in metropolitan Lima in 2018 (Bulnes-Bedón, Alvarez-Taco, & Morales-Isasi, 2018). The degree of anxiety was evaluated using Zung's 20-item Self-rating Anxiety Scale (Dunstan & Scott, 2020). In its Spanish version, each item was evaluated on a Likert scale, rated from 1 to 4, with the lowest score being 20 and the highest 80. Scores 50-59, 60-69, and 70 and over represent mild, moderate, and severe anxiety, respectively.

Procedure

Once permission had been granted by the universities and the study had been approved by the ethics committee of Antenor Orrego Private University, data were collected in March and April of 2022. To corroborate their status as registered students, standardized messages were sent with an invitation to participate in the study to the institutional emails of medical students from UPAO, UCV, and UNP universities. If the students agreed to participate, they were contacted via WhatsApp on their personal cell phones to explain the objective of the study, the selection criteria, the informed consent form and the way the measurement tool

Table 1
Anxiety level and psychosomatic features of medical students from three university medical faculties in Northern Peru

	Psychosomatic features				p-value
	Absent n (%)	Mild n (%)	Moderate n (%)	Severe n (%)	
Anxiety level	77 (21.9)	126 (35.8)	105 (29.8)	44 (12.5)	
Absent	73 (94.8)	116 (92.0)	76 (72.3)	34 (77.3)	
Mild	4 (5.2)	9 (7.1)	28 (26.7)	8 (18.1)	< .001
Moderate	0 (0)	1 (.9)	1 (1)	2 (4.6)	

Source: UPAO, UCV and UNP student data sheets.

should be completed. Students who met the selection criteria and voluntarily agreed to participate were sent the online link to Google Forms.

Statistical analysis

Data from the surveys undertaken were entered into a Microsoft Excel spreadsheet for cleaning and purging. STATA version 16 was used for the analysis. The PHQ-15 screening instrument and Zung's test scores were categorized and regarded as qualitative variables, which were presented in frequency and percentage. The quantitative variable age was presented as a mean and standard deviation. The chi-square test was used to find an association. The PRc and PRa were subsequently calculated (adjusting for age, gender, study cycle and grade retention).

Ethical considerations

There were no risks or benefits for any participant. Answering the virtual survey did not involve any expense, nor were financial incentives provided. Data confidentiality was maintained, and participant anonymity respected. The project was approved by the ethics committee of Antenor Orrego Private University with code RR N° 3335-2016-R-UPAO.

RESULTS

Association between anxiety and psychosomatic features

As can be seen from Table 1, the degrees of PF were categorized, determining the level of anxiety for each degree. It was found that over 75% of the students surveyed presented PF. The maximum level of anxiety found was moderate, with a progressive increase being observed in cases with the most severe degrees of PF. The association was highly significant.

Factors associated with psychosomatic features

Table 2 shows that being female increases the risk of presenting psychosomatic features by 45% compared to males. Presenting a mild and moderate level of anxiety increases the risk of PF by 40% and 70% respectively. There were no significant associations with age, studying clinical sciences or grade retention.

DISCUSSION AND CONCLUSION

This study, which was designed to determine the association between psychosomatic features and anxiety levels in university students, included a total of 352 students.

It was found that 35.8% presented mild, 29.8% moderate, and 12.5% severe psychosomatic features, with a total of 78.1% presenting some type of psychosomatic feature. Despite these figures, there is a dearth of information in the recent literature on the prevalence of psychosomatic features in university students. However, this is associated with the high prevalence of psychosomatic features in health workers, which is above 75% (López-Rodríguez, García-Gracia, Ponce-Martín, Arranz-Ballesteros, & Parejo-Aguilera, 2021; Uribe-Prado, 2020).

The degree of anxiety found was distributed as follows: 13.9% presented a mild degree of anxiety and 1.1% a mod-

Table 2
Bivariate and multivariate analysis of psychosomatic features and their association with the variables studied

	n (%)	Bivariate PRc (95% CI)	p-value	Multivariate PRa (95% CI)	p-value
Female sex	199 (56.5)	1.5 (1.27, 1.78)	< .001	1.45 (1.23, 1.71)	< .001
Age (M ± SD)	23.2 ± 3.7	.98 (.96, 1.00)	.147	.98 (.96, 1.01)	.427
Clinical sciences (≥ 6th cycle)	220 (62.4)	1.08 (.91, 1.27)	.343	1.11 (.92, 1.33)	.24
Grade retention	172 (48.9)	.91 (.78, 1.05)	.229	.91 (.78, 2.34)	.22
Anxiety level		1.24 (1.26, 1.61)	< .001		
Absent	299 (84.9)			References	
Mild	49 (13.9)			1.4 (1.20, 1.63)	< .001
Moderate	4 (1.1)			1.7 (1.24, 2.34)	.001

Source: UPAO, UCV and UNP student data sheets.

erate degree. The degree of anxiety found was distributed as follows: 13.9% showed a mild and 1.1% a moderate degree. These results vary according to the type of test used to determine the degree of anxiety. Thus, for example, a study conducted by [Ordóñez-Galeano \(2020\)](#) in Guatemala using Beck's inventory with first and last year medical students found that 68% and 66% experienced mild anxiety, and 41% and 57% moderate anxiety respectively ([Ordóñez-Galeano, 2020](#)). At the international level, in Spain, [Gutiérrez-Pastor I.](#) used Goldberg's Anxiety and Depression Scale (GADS) for medical students at a Spanish university, finding a global anxiety rate of 54.9% ([Gutiérrez-Pastor, Quesada-Rico, Gutiérrez-Pastor, Nouni-García, and Carratalá-Munuera, 2021](#)). Peruvian studies have revealed a high incidence of anxiety, ranging from 23% according to the Beck Inventory to 75.4% using the Generalized Anxiety Disorder Assessment (GAD-7; [Armas-Elguera, Talavera, Cárdenas, & de la Cruz-Vargas, 2021](#); [Saravia-Bartra, Cazorla-Saravia, & Cerdillo-Ramirez, 2020](#)). These values can be influenced by social and political settings such as the COVID-19 Pandemic ([Saravia-Bartra et al., 2020](#)) as well as by the year of study and the consumption of harmful products such as alcohol and tobacco ([Robles-Mariños, Angeles, & Alvarado, 2022](#)).

Our study found a low probability of the association between the degree of anxiety and psychosomatic features being due to chance ($p < .001$). It also discovered that the presence of any degree of anxiety increased by 40% (PRC = 1.4, 95% CI [1.26, 1.61], p value $\leq .001$) the possibility of presenting psychosomatic features compared to students who had no anxiety. In the statistical analysis adjusted by age, gender, year of study and grade retention, it was found that female students were 45% more likely (RPa = 1.45, CI 95% [1.23, 1.73], p value $\leq .001$) to present psychosomatic features than male students. In addition, having a mild or moderate level of anxiety meant that students were 40% (PRa = 1.4, 95% CI [1.20, 1.63], p value $\leq .001$) and 70% (PRa = 1.7, 95% CI [1.24, 2.34], p value = .001) more likely to present psychosomatic features respectively, compared with those who do not experience anxiety. This can be explained by the nature of these somatic symptoms, which are often considered to be vague somatoform disorders superimposed on depression or anxiety or organic causal attributions. This means that the presence of a mental condition is necessary for the presence of a psychosomatic symptom ([Trebin, 2020](#)). These stressors produce selective alterations of large-scale brain networks involved in cognitive control and the regulation, and processing of emotions, stress, and somatic-visceral perception, leading to various psychosomatic features such as headaches, low back pain, joint pain, retro-ocular pain, nausea, dizziness, vomiting, seizures (conversion syndrome), and fainting ([Rossetti et al., 2021](#)).

The study concludes that psychosomatic features are found in approximately 75% of medical students and anxiety in approximately 15%. Mild and moderate anxiety is an

influential factor in psychosomatic features, as is being female. Given this situation, periodic evaluations and anxiety management sessions could be proposed to reduce psychosomatic features to ensure the mental health of future doctors, which will have a positive effect on their academic training.

Funding

The present study was funded by the authors.

Conflict of interest

The authors declare that they have no conflicts of interest.

Acknowledgements

We would like to thank the university chancellors for their support.

REFERENCES

- Alzahrani, A., Alghamdi, A., Alqarni, T., Alshareef, R., & Alzahrani, A. (2019). Prevalence and predictors of depression, anxiety, and stress symptoms among patients with type II diabetes attending primary healthcare centers in the western region of Saudi Arabia: a cross-sectional study. *International Journal of Mental Health Systems, 13*, 48. doi: 10.1186/s13033-019-0307-6
- Anton, N. E., Rendina, M. A., Hennings, J. M., Stambro, R., Stanton-Maxey, K. J., & Stefanidis, D. (2021). Association of Medical Students' Stress and Coping Skills With Simulation Performance. *Simulation in Healthcare, 16*(5), 327-333. doi: 10.1097/sih.0000000000000511
- Armas-Elguera, F., Talavera, J. E., Cárdenas, M. M., & de la Cruz-Vargas, J. A. (2021). Trastornos del sueño y ansiedad de estudiantes de Medicina del primer y último año en Lima, Perú. *FEM: Revista de la Fundación Educación Médica, 24*(3), 133-138. doi: 10.33588/fem.243.1125
- Atta, I. S., & Almilaibary, A. (2022). The Prevalence of Stress Among Medical Students Studying an Integrative Curriculum During the COVID-19 Pandemic. *Advances in Medical Education and Practice, 13*, 35-45. doi: 10.2147/amep.s345330
- Bartmann, C., Fischer, L. M., Hübner, T., Müller-Reiter, M., Wöckel, A., McNeill, R. V., ... Diessner, J. (2021). The effects of the COVID-19 pandemic on psychological stress in breast cancer patients. *BMC Cancer, 21*(1), 1356. doi: 10.1186/s12885-021-09012-y
- Bulnes-Bedón, M. S., Alvarez-Taco, C. L., & Morales-Isasi C. (2018). Ansiedad social y regulación emocional en adolescentes de Lima metropolitana con y sin presencia de síntomas somáticos. *Temática Psicológica, 14*(14), 51-68. doi: 10.33539/tematpsicol.2018.n14.1810
- Cabanach, R. G., Souto-Gestal, A., & Fernández Cervantes, R. (2017). Perfiles de regulación emocional y estrés académico en estudiantes de fisioterapia. *European Journal of Education and Psychology, 10*(2), 57-67. doi: 10.1016/j.ejeps.2017.07.002
- de La Rosa-Rojas, G., Chang-Grozo, S., Delgado-Flores L., Oliveros-Lijap L., Murillo-Pérez D., Ortiz-Lozada R., ... Carreazo, N. Y. (2015). Niveles de estrés y formas de afrontamiento en estudiantes de Medicina en comparación con estudiantes de otras escuelas; lima: Universidad Peruana de Ciencias Aplicadas; 2015. *Gaceta Médica de México, 151*(4), 443-449.
- Dunstan, D. A., & Scott, N. (2020). Norms for Zung's Self-rating Anxiety Scale. *BMC Psychiatry, 20*(1), 90. doi: 10.1186/s12888-019-2427-6
- Galiano-Ramírez, M. de la C., Castellanos-Luna, T., & Moreno-Mora, T. (2016). Manifestaciones somáticas en un grupo de adolescentes con ansiedad. *Revista Cubana de Pediatría, 88*(2), 195-204.
- Gutiérrez-Pastor, I., Quesada-Rico, J. A., Gutiérrez-Pastor, A., Nouni-García, R., & Carratalá-Munuera, M. C. (2021). Depresión, ansiedad y salud autopercibida en estudiantes de Medicina: un estudio transversal. *Revista Española de Educación Médica, 2*(2), 21-31.
- Kroenke, K., Spitzer, R. L., & Williams, J. B. (2002). The PHQ-15: validity of a new measure for evaluating the severity of somatic symptoms. *Psychosomatic Medicine, 64*(2), 258-266. doi: 10.1097/00006842-200203000-00008

- López-Rodríguez, B., García-Gracia, E., Ponce-Martín, P., Arranz-Ballesteros, B., & Parejo-Aguilera, M. C. (2021). Entorno psicosocial y manifestaciones psicósomáticas en enfermeras de una unidad de quirófano. *Revista de la Asociación Española de Enfermería Quirúrgica*, 46(1885-2548), 21-27.
- Maria-Jereyard, K. G., & Bhuvaneswari, U. (2019). Psychosomatic symptoms and Higher Secondary School students. *Indian Journal of Applied Research*, 9(11).
- Nochaiwong, S., Ruengorn, C., Thavorn, K., Hutton, B., Awiphan, R., Phosuya, C., ... Wongpakaran, T. (2021). Global prevalence of mental health issues among the general population during the coronavirus disease-2019 pandemic: a systematic review and meta-analysis. *Scientific Reports*, 11(1), 10173. doi: 10.1038/s41598-021-89700-8
- Ordóñez-Galeano, R. A. (2020). Depresión y ansiedad en estudiantes de medicina. *Revista Ciencia Multidisciplinaria CUNORI*, 4(2), 15-21. doi: 10.36314/cunori.v4i2.123
- Quito-Calle, J. V., Tamayo-Piedra, M. del C., Buñay-Barahona, D. P., & Neira-Cardenas, O. S. (2017). Estrés académico en estudiantes de tercero de bachillerato de unidades educativas particulares del Ecuador. *Revista Electrónica de Psicología Iztacala*, 20(3), 253-276.
- Robles-Mariños, R., Angeles, A. I., & Alvarado, G. F. (2022). Factores asociados con la ansiedad por la salud en estudiantes de Medicina de una universidad privada en Lima, Perú. *Revista Colombiana de Psiquiatría*, 51(2), 89-98. doi: 10.1016/j.rcp.2020.11.002
- Rossetti, M. G., Delvecchio, G., Calati, R., Perlini, C., Bellani, M., & Brambilla, P. (2021). Structural neuroimaging of somatoform disorders: A systematic review. *Neuroscience & Biobehavioral Reviews*, 122, 66-78. doi: 10.1016/j.neubiorev.2020.12.017
- Sajid, A., Ahmad, T., & Khalid, T. (2015). Stress in medical undergraduates; its association with academic performance. *Bangladesh Journal of Medical Science*, 14(2), 135-141. doi: 10.3329/bjms.v14i2.21815
- Salam, A., Mahadevan, R., Abdul Rahman, A., Abdullah, N., Abd Harith, A. A., & Shan, C. P. (2015). Stress among First and Third Year Medical Students at University Kebangsaan Malaysia. *Pakistan Journal of Medical Sciences*, 31(1), 169-173. doi: 10.12669/pjms.311.6473
- Saravia-Bartra, M. M., Cazorla-Saravia, P., & Cedillo-Ramirez, L. (2020). Nivel de ansiedad de estudiantes de medicina de primer año de una universidad privada del Perú en tiempos de Covid-19. *Revista de la Facultad de Medicina Humana*, 20(4), 568-573. doi: 10.25176/rfmh.v20i4.3198
- Tollos, I., Theodorakopoulou, A., & Christodoulou, G. N. (2021). Stress and pathophysiological mechanisms for the development of psychosomatic disease. *Psychiatriki*, 32(2), 148-156. doi: 10.22365/jpsych.2021.023
- Trebin, E. (2020). Psychosocial and Somatoform Disorders. *Deutsches Ärzteblatt International*, 116(8), 134. doi: 10.3238/arztebl.2020.0134a
- Uribe-Prado, J. F. (2020). Riesgos psicosociales, burnout y factores psicósomáticos en servidores públicos. *Investigación Administrativa*, 49, 125. doi: 10.35426/iav49n125.03