

# Clinical practice stressors and anxiety in nursing students during COVID-19

Nancy Griselda Pérez Briones,<sup>1</sup> Karla Rosales Vázquez,<sup>1</sup> Braulio Josué Lara Reyes,<sup>1</sup> Dafne Astrid Gómez Melasio,<sup>1</sup> Anahí Ruiz Lara,<sup>2</sup> Pedro Enrique Trujillo Hernández<sup>1</sup>

<sup>1</sup> Facultad de Enfermería "Dr. Santiago Valdés Galindo", Universidad Autónoma de Coahuila, Saltillo, México.

<sup>2</sup> Escuela de Ciencias de la Salud, Unidad Norte, Universidad Autónoma de Coahuila, Piedras Negras, México.

## Correspondence:

Dafne Astrid Gómez Melasio  
Facultad de Enfermería "Dr. Santiago Valdés Galindo", Universidad Autónoma de Coahuila,  
Calz. Francisco I. Madero 1237,  
Zona Centro,  
25000 Saltillo, Coah., México.  
Phone: +52 (844) 414-2588  
Email: [dafne.gomez@uadec.edu.mx](mailto:dafne.gomez@uadec.edu.mx)

Received: 7 November 2022

Accepted: 21 February 2023

## Citation:

Pérez Briones, N. G., Rosales Vázquez, K., Lara Reyes, B. J., Gómez Melasio, D. A., Ruiz Lara, A., & Trujillo Hernández, P. E. (2023). Clinical practice stressors and anxiety in nursing students during COVID-19. *Salud Mental*, 46(6), 287-293.

DOI: 10.17711/SM.0185-3325.2023.037



## ABSTRACT

**Introduction.** The suspension of clinical practices in the nursing profession during COVID-19 generated a decrease in the students' abilities and skills which made them vulnerable to more frequent to clinical practice stressors and levels of anxiety. **Objective.** To determine the relation between clinical practice stressors and anxiety levels in nursing students during COVID-19. **Method.** A descriptive correlational study with convenience sampling was carried out in 169 students enrolled from third to eighth semesters in a public university in the city of Saltillo, Coahuila. For the measurement of variables a sociodemographic data card, the KEZKAK Questionnaire in its Spanish version to measure the clinical practice stressors and the Beck Anxiety Inventory were used. Data collection was through a digital link during August-December 2021. Descriptive and inferential statistics were used through SPSS V.26. **Results.** 57.4% reported a medium level of stress and 62.7% a low level of anxiety; however, there are stressors from clinical practice that lead to increased anxiety. A statistically significant positive correlation was found between the study variables ( $r = .411, p < .001$ ). **Discussion and conclusion.** The challenges in clinical practice experienced during COVID-19 highlight that generations of future nursing professionals face a scenario that is the product of a post-pandemic era, so it is necessary to strengthen their knowledge and confidence to motivate them to feel safe within the hospital context.

**Keywords:** stressors, anxiety, nursing students, coronavirus infections.

## RESUMEN

**Introducción.** La suspensión en las prácticas clínicas en la profesión de enfermería durante la COVID-19, generó en los estudiantes disminución de sus habilidades y destrezas que los hizo vulnerables a presentar con mayor frecuencia estresores de la práctica clínica y niveles de ansiedad. **Objetivo.** Determinar la relación entre los estresores de la práctica clínica y los niveles de ansiedad en estudiantes de enfermería durante la COVID-19. **Método.** Se realizó un estudio de tipo descriptivo correlacional con muestreo por conveniencia en 169 estudiantes de tercero a octavo semestre matriculados en una universidad pública de la ciudad de Saltillo Coahuila. Para la medición de variables se utilizó la cédula de datos sociodemográficos, el Cuestionario KEZKAK para medir los estresores de la práctica clínica y el Inventario de Ansiedad de Beck. La colecta de datos fue a través de un enlace digital durante agosto-diciembre 2021. Se utilizó estadística descriptiva e inferencial a través del SPSS V.26. **Resultados.** El 57.4% refiere un nivel medio de estrés y el 62.7% un bajo nivel de ansiedad; sin embargo, existen estresores provenientes de la práctica clínica que propician el aumento de ansiedad. Se encontró una correlación positiva estadísticamente significativa entre las variables de estudio ( $r = .411, p < .001$ ). **Discusión y conclusión.** Los retos en la práctica clínica experimentados durante la COVID-19, resaltan que las generaciones de futuros profesionistas en enfermería se enfrentan a un panorama producto de una época post-pandemia, por lo que es necesario afianzar sus conocimientos y su confianza para motivarlos a sentirse seguros dentro del contexto hospitalario.

**Palabras clave:** estresores, ansiedad, estudiantes de enfermería, infecciones por coronavirus.

## INTRODUCTION

The university stage is the most important period of time in the lives of students, because it represents the consolidation of their life project. In the nursing training process, practice and theory complement each other in order to bring the student closer to the healthcare reality and from this, they acquire professional skills suitable for their insertion in the workplace (Cardona-Arias, Pérez-Restrepo, Rivera-Ocampo, Gómez-Martínez, & Reyes, 2015; Lara, 2020).

As far as clinical practice is concerned, it is a process that involves knowledge, they acquire professional skills suitable for their insertion in the workplace. Abilities, skills, attitudes, and values that are important for carrying out procedures. As well as, for the acquisition of experience in the hospital environment, which contributes to providing better care to patients (Tiga, Parra, & Domínguez, 2014).

Given the care demands during clinical practice, it has been possible to identify how nursing students perceive stressful situations during their internship period, derived from the nurse-patient relationship and the health team, making them vulnerable to presenting physical reactions such as insomnia, fatigue, muscle tension, among other symptoms. Likewise, they refer the presence of psychological symptoms, such as stress, anxiety, alterations in the sleep-wake state, inability to concentrate, and relax. Del Risco, Gómez, and Hurtado (2015) mentions that the main sources of stress of the student population in training come from situations in the clinical area due to contact with diseases, pain, suffering, death, cardiorespiratory arrest, and the difference between practice and theory (Moya et al., 2013; Villanueva & Ugarte, 2017).

On the other hand, the emergence of a new virus called SARS-CoV-2 or COVID-19 disease in China at the end of 2019 caused a general health problem that affected the world population. Its rapid dispersion caused the establishment of the pandemic situation to be declared in March 2020, forcing governments to take containment measures in all social areas, including the educational sector (Barros et al., 2020; Caldera-Villalobos et al., 2020; Chih-Cheng, Tzu-Ping, Wen-Chien, Hung-Jen, & Po-Ren, 2020).

Therefore, the governments opted for the suspension of face-to-face class activities, including clinical practices, in compliance with the recommendations of the health sector. The most affected are those students in the health area, especially nursing, where the student reinforces knowledge by developing procedures in the hospital environment and during clinical simulation environments of the institutions (Alemán, Vera, & Patiño-Torres, 2020).

The adequacy of the restrictions in the nursing educational system influenced the psychological well-being of the students when they returned to clinical practice, because for more than a year their source of learning was only through online classes, causing insecurity in the procedure

performance and in relationships with other members of the hospital environment (Sánchez & Talavera, 2021).

Nursing students are prone to clinical practice stressors derived from situations within the hospital that force them to make effective, efficient, safe, and up-to-date decisions (Rabelo-Silva, Monteiro, & Lumertz, 2022). It should be noted that the confinement generated psychosocial pressure as a result of the online classes and the lack of the clinical practice period. Therefore, symptoms related to stress and anxiety were presented, as a result of the constant concern of not having completed their internship period or having performed the procedures in the clinical simulation center during the start of the COVID-19 pandemic.

Currently, it is imperative for the nursing profession to combat stress and anxiety because they interfere with the quality of patient care. Therefore, nursing teachers must teach strategies that help reduce stress and anxiety, as well as positively support students at the individual and family level in order to increase their confidence, security, skills and abilities skills within the hospital (Mamani & Mamani, 2022; Valero, Vélez, Duran, & Torres, 2020). For all of the above, the objective of this research is to determine the relationship between the clinical practice stressors and the levels of anxiety in nursing students during COVID-19.

## METHOD

### Design of the study/Participants

A quantitative descriptive correlational study was carried out with non-probabilistic convenience sampling in 169 nursing students over 18 years of age, without psychological and/or pharmacological treatment, enrolled from the third to the eighth semester at a public university in the city of Saltillo, Coahuila.

### Procedure

The data collection took place during August-December 2021 through a digital link that included informed consent and measurement instruments, which was shared by WhatsApp with the group leaders of the semesters to be investigated.

### Measurements

To obtain the data, the following instruments were used, which were transcribed on the Microsoft Forms platform:

- Sociodemographic data card: It was prepared to characterize the study population, which included sex, age, marital status, and current semester.
- KEZKAK Bilingual Questionnaire: It was used to measure the clinical practice stressors, which obtained a Cronbach's Alpha of .96. Likewise, it consists of nine dimensions and 41 items, with four

Likert-type response options ranging from 0: Not at all, 1: Somewhat, 2: Quite a bit, and 3: A lot. Each dimension is measured according to the following cut-off points established by Madeley (2020):

- Dimension 1-Lack of competence (items 1,2,3,4,5,6,13,15,16,17,26): low (0-11 points), medium (12-22 points), high (23-33 points).
  - Dimension 2-Contact with suffering (items 9, 10,14,18,27,29,31,32,34,39): low (0-11 points), medium (12-22 points), high (21-30 points).
  - Dimension 3-Relationship with tutor and classmates (items 1,12,19,20,25,28): low (0-6 points), medium (7-12 points), high (13-18 points).
  - Dimension 4-Powerlessness and uncertainty (items 2,3,6,14,17,20,23,32,36,38,41): low (0-11 points), medium (7-12 points), high (23-33 points).
  - Dimension 5-Not controlling the relationship with the patient (items 5,7,17,20,29,30,33,39): low (0-8 points), medium (9-16 points), high (17-24 points).
  - Dimension 6-Emotional involvement (items 8,21,22,31): low (0-4 points), medium (5-8 points), high (9-12 points).
  - Dimension 7-Damaging in relation to the patient (items 1,14,15,24,26): low (0-5 points), medium (6-10 points), high (11-15 points).
  - Dimension 8-The patient seeks an intimate relationship (items 37,40): low (0-2 points), medium (3-4 points), high (5-6 points).
  - Dimension 9-Overload (items 30,31,34,35,36): low (0-5 points), medium (6-10 points), high (11-15 points).
- Beck Anxiety Inventory (BAI): It was used to determine the anxiety levels of the student population. The questionnaire showed a Cronbach's Alpha of .92 and is made up of 21 items with four Likert-type response options ranging from 0, which represents "nothing" or no anxiety, to 3, which represents "severely" or the level of anxiety. The cut-off points are: very low anxiety: 0-21 points; moderate anxiety: 22-35; and severe anxiety: > 36 points.

### Statistical analysis

For data processing, the statistical program Statistical Package for the Social Sciences (SPSS) Version 26 for Windows was used. Once the data collection was completed, the database was created. Descriptive statistics were applied through frequencies and percentages, as well as measures of central tendency and dispersion.

The consistency of the instruments was determined through Cronbach's Alpha coefficient for Likert-type an-

swers. As well as, a goodness or fit analysis was performed with the Kolmogorov-Smirnov test, a curve with normal distribution was observed ( $p > .05$ ). Therefore, Pearson's parametric correlation model was used.

### Ethical considerations

The present study was approved by the Research Ethics Committee of the Faculty of Nursing "Dr. Santiago Valdés Galindo" of the Universidad Autónoma de Coahuila (FAENUS-CEI-EXT-2021-08) and was based on Regulations of the General Health Law on Health Research, considering it a minimal risk investigation since the data recording was carried out through common procedures, such as the application of a digital link. Likewise, it adhered to the international guidelines of the Helsinki Code to protect the confidentiality of the participants. In addition, to respect the freedom to withdraw their consent and stop participating in the study at the time they decided.

### RESULTS

A total of 207 participants were collected, of which 169 met the inclusion criteria, there was a predominance of the female population occupying 85.5% of the total number of respondents, with an average age of 20 years ( $\bar{x} = 20.20$ ,  $SD = 1,481$ ) with ages between 18 and 27 years. Regarding the current academic semester, a mean of 5.15 and a standard deviation of 1.239 were obtained; obtaining that the fifth semester students were the ones that showed a greater participation. Regarding marital status, 92.9% are single, 4.1% are married, and the remaining 3% are in a cohabitation.

To measure the stress produced by the clinical practice stressors, the total sum of the items was carried out to later classify the data into three low (25), medium (50), and high (75) percentiles, finding that more than 50% of the participants report a medium level of stress and 15.4% show a high level (Table 1).

On the other hand, the KEZKAK Questionnaire has nine additional dimensions to the level of general perceived stress. Each of these dimensions has measurement levels, which mostly came out at a medium level, corroborating the level of stress perceived by the participants in clinical practice without there being any particular alteration. It should

Table 1  
Stress level in nursing students

Stress level	f	%
Low	46	27.2
Medium	97	57.4
High	26	15.4

Note: f = frequency, % = percentage, n = 169

Table 2  
Clinical practice stressors by dimension in nursing students

Dimensions	Low		Medium		High	
	f	%	f	%	f	%
Dimension 1: Lack of competence	64	37.9	67	39.6	38	22.5
Dimension 2: Contact with suffering	64	37.9	84	49.7	21	12.4
Dimension 3: Relationship with tutor and classmates	64	37.9	89	52.7	16	9.5
Dimension 4: Powerlessness and uncertainty	36	21.3	95	56.2	38	22.5
Dimension 5: Not controlling the relationship with the patient	49	29	89	52.7	31	18.3
Dimension 6: Emotional involvement	83	49.1	70	41.4	16	9.5
Dimension 7: Damaging in relation to the patient	63	37.3	77	45.6	29	17.2
Dimension 8: The patient seeks an intimate relationship	50	29.6	54	32	65	38.5
Dimension 9: Overload	60	35.5	85	50.3	24	14.2

Note: f = frequency, % = percentage, n = 169

be noted that the most prevalent were dimension 3: “Relationship with the tutor and/or colleagues” and dimension 5: Not controlling the relationship with the patient with 52.7% and dimension 4: “importance and uncertainty” with 56.2% (Table 2).

Regarding the stressors that cause between a lot and a lot of stress, the students refer to more than 30% that pricking themselves with an infected needle causes them stress; being wrong; not knowing what to do; being in an emergency; the differences between what is learned in theory and reality when executed in practice. As well as, 46.7% are stressed not to find the doctor when needed and that the patient touches some part of her body (Table 3).

Regarding the anxiety variable, it was found that only 27.2% show a moderate level, while 62.7% of those surveyed show a very low level of anxiety. Of the 21 items that comprise the BAI, a significant increase is observed in 3 items related to anxious symptomatology, which are “Nervous” with 37.3%, “With digestive problems” with 24.9%, and “Being unable to relax” with 30.2% (Table 4).

Likewise, a moderate positive relationship with statistical significance was found between clinical practice stressors and anxiety levels ( $r = .411$ ,  $p < .001$ ), which indicates that the greater the presence of clinical practice stressors higher clinical level of anxiety perceived by participants in the hospital environment (Table 5).

In summary, the results obtained show that the level of anxiety is perceived as very low by the participants. However, there are stressors of clinical practice that favor the presence of stress and therefore an increase in anxiety among students when executing procedures due to lack of practice within the hospital environment during COVID-19.

## DISCUSSION AND CONCLUSION

The sociodemographic profile showed that the female population predominates in nursing, agreeing with [Castillo, Chacón, and Díaz-Véliz \(2016\)](#) who found that women made up the majority of their total population (64.1%), the average age they acquired was 19 years ( $ED = .1$ ), which coincided with this investigation, where a mean age of 20 years was found ( $ED = 1.481$ ). It should be noted that, in the usual context of studies within the nursing literature, the female sex has predominated ([Bodys-Cupak, Majda, Skowron, Zalewska-Puchala, & Trzcinka, 2018](#); [Hernández, 2020](#)). The results reported by [López, Orbañanos, and Cibanal \(2013\)](#) in the study they carried out, with the aim of identifying stressors in clinical practices and their evolution, were not representative or generalizable to the male sex.

Historically, the nursing profession has been exercised mostly by women, however, there may be variations between the different sociocultural contexts, in addition, currently the number of men who practice the nursing profession is increasing. Research has even been carried out that includes only male nursing students ([Mahadeen, Abushaikha, & Habshneh, 2017](#)).

Regarding the stressors of clinical practice, a higher prevalence was found in dimension 4: “Helplessness and Uncertainty” (56.2%); the most prominent items being: “doing my job poorly and harming the patient” and “screwing up”. The previous findings agree with what was stated by [López et al. \(2013\)](#), [Fandiño, Henríquez, and Rivera, \(2020\)](#), and [Velásquez-Muñoz, Cornejo-Aravena, Bustamante-Toncio, and Mella-Guzmán \(2018\)](#). [Fandiño et al., \(2020\)](#) found that the participating women obtained higher scores in the intermediate to high values (somewhat - quite) of the items of the “Helplessness and Uncertainty” dimension, compared to men.

**Tabla 3**  
*Clinical practice stressors with greater frequency in nursing students*

Clinical practice stressors		f	%
Prick me with an infected needle	Not at all	27	16
	Somewhat	39	23.1
	Quite a bit	49	29
	A lot	32	<b>32</b>
Confuse the medication	Not at all	27	16
	Somewhat	35	20.7
	Quite a bit	48	28.4
	A lot	59	<b>34.9</b>
"Mess up" or make a mistake	Not at all	13	7.7
	Somewhat	35	20.7
	Quite a bit	59	34.9
	A lot	62	<b>36.7</b>
Find myself in a situation without knowing what to do	Not at all	12	7.1
	Somewhat	42	24.9
	Quite a bit	58	<b>34.3</b>
	A lot	57	33.7
Find myself in an emergency situation	Not at all	19	11.2
	Somewhat	58	34.3
	Quite a bit	61	<b>36.1</b>
	A lot	31	18.3
Not finding the doctor when the situation requires it	Not at all	16	9.5
	Somewhat	35	21.3
	Quite a bit	79	<b>46.7</b>
	A lot	38	22.5
That the patient touches certain parts of my body	Not at all	23	13.6
	Somewhat	25	14.8
	Quite a bit	42	24.9
	A lot	79	<b>46.7</b>
Differences between what we learned in class and what we see in practice	Not at all	13	7.7
	Somewhat	43	25.4
	Quite a bit	62	<b>36.7</b>
	A lot	51	30.2

Note: f = frequency, % = percentage, n = 169

López et al. (2013) mention that Dimension 4: "Impotence and Uncertainty" can be altered in students by the desire to help the patient, seeking to be better in the tasks they perform, however, not being able to satisfy the needs of the patients or the correct execution of a procedure, can be frustrating for the student. The insecurity in students of not performing the procedures correctly could have been considerably increased by the suspension of clinical practice during the pandemic, considering that they did not acquire

**Table 4**  
*Anxious symptomatology frequently presented in nursing students*

Anxious symptomatology		f	%
Unable to relax	Very low anxiety	41	24.3
	Low anxiety	65	<b>38.5</b>
	Moderate anxiety	51	30.2
	Severe anxiety	12	7.1
Fearing the worst will happen	Very low anxiety	35	20.7
	Low anxiety	69	<b>40.8</b>
	Moderate anxiety	47	27.8
	Severe anxiety	18	10.7
Accelerated heartbeat	Very low anxiety	69	<b>40.8</b>
	Low anxiety	55	32.5
	Moderate anxiety	36	21.3
	Severe anxiety	9	5.3
Nervous	Very low anxiety	21	12.4
	Low anxiety	56	33.1
	Moderate anxiety	63	<b>37.3</b>
	Severe anxiety	29	17.2
Digestive problems	Very low anxiety	65	<b>38.5</b>
	Low anxiety	52	30.8
	Moderate anxiety	42	24.9
	Severe anxiety	10	5.9

Note: f = frequency, % = percentage, n = 169

sufficient skills to carry out the procedures. It is expected that, with the passage of time, the reintegration into the hospital context and the experience acquired, will increase the confidence and security of the students and with this, they will be able to face stressful situations in a positive way.

Analyzing each of the items that make up the KEZKAK Inventory, significant increases were observed in the frequencies of the items "Confusing myself with the medication", "The difference between theory and practice", "Puncture myself with an infected needle", "Find myself in an emergency situation" and "Receive contradictory orders", these findings

**Table 5**  
*Clinical practice stressors related to anxiety levels in nursing students*

Variables	1	2
1. Clinical practice stressors	1	.411 .001**
2. Anxiety	.411 .001**	1

Note: \*\*Correlation is significant at the .001 level, n = 169



are similar to those reported by Arias, Montoya, Villegas, and Rodríguez-Gázquez (2018), Fandiño et al. (2020), and Velásquez-Muñoz et al. (2018). Arias et al. (2018) found as main stressors, feeling that they are doing their job badly, harming the patient, doing physical harm to the patient and getting confused in the medication. Velásquez-Muñoz et al. (2018) identified that the highest prevalence of stressors in students were those related to competencies, impotence, and uncertainty in performing the procedures.

Sánchez (2019) identified that situations such as those mentioned can be determined as causes of stress, due to the responsibility they imply for the student and the negative repercussions that some of them could have on the patient, which may occur more frequently as consequence of the return to hospital clinical practice after the pandemic.

In the present investigation, a higher frequency was found in the item "Let the patient touch certain parts of my body", which was not found as a relevant finding in the literature review carried out, since it was one of the least valued items of the Inventory. This could be due to the cultural context of gender violence in which Mexico finds itself, and that a large part of the sample for this research were women. Likewise, it is considered that the presence of this stressor is a consequence of the fear of avoiding the spread of the SARS-CoV-2 virus.

Regarding the anxiety variable, the results indicate that the majority of the population surveyed perceive low levels (62.7%), which is consistent with the results provided by Clemente (2021) and Fandiño et al. (2020). However, there are studies in which anxiety levels are measured with the State-Trait Anxiety Inventory (STAI), which assesses the individual's personal perception of their anxiety through subjective feelings, unlike the Beck Anxiety Inventory (BAI), which evaluates the somatic symptoms experienced before the stressor in question. Therefore, the results through the BAI are useful to show the most relevant symptoms of anxiety. In this study, the items "Unable to relax," "Nervous," and "With digestive problems" turned out to be the most prevalent, being one of the most common reactions presented in a picture of anxiety, similar to what was obtained in Fandiño et al. (2020).

However, despite the low levels of anxiety perceived by the student population, a significant relationship was found between clinical practice stressors and anxiety levels. This agrees with the results found by Clemente (2021) and Hernández (2020) who also obtained a positive and significant relationship. Hernández (2020) evaluated the levels of stress and anxiety during clinical practice, finding moderate levels during the course, identifying the lack of competence and impotence and uncertainty as the main stressors that cause the increase in the prevalence of anxiety in students.

The findings of this study show a positive correlation between the stressors of clinical practice and the levels of anxiety manifested, so it can be identified that the challenges rep-

resented by the hospital environment have an impact on the psychosocial health of students. However, it is not possible to determine the impact that these stressors have had on their performance. Despite this, it was possible to identify some of the situations that cause the most stress, which are consistent with the findings of different authors in the literature.

It should be noted that the new generations of future nursing professionals who resume their clinical practices face a different scenario as a result of a post-pandemic era, so the results must be taken into account to improve the education of students leaving online education and face the real situation, in order not only to increase their abilities and/or skills, but also to strengthen, motivate, and give them the necessary confidence to feel safe in the hospital context, in order to train future quality professionals.

### Funding

None.

### Conflict of interest

The authors declare they have no conflicts of interest.

### Acknowledgements

To the Universidad Autónoma de Coahuila for allowing the project to be carried out, as well as the study subjects for their willingness to contribute to the investigation.

### REFERENCES

- Alemán, I., Vera, E., & Patiño-Torres, M. J. (2020). COVID-19 y la educación médica: retos y oportunidades en Venezuela. *Educación Médica*, 21(4), 272-276. doi: 10.1016/j.edumed.2020.06.005
- Arias, L. Y., Montoya, L. I., Villegas, A. F., & Rodríguez-Gázquez, M. de los A. (2018). Estresores en las prácticas clínicas de los estudiantes de enfermería de una universidad pública en Colombia. *Investigación en Enfermería: Imagen y Desarrollo*, 20(1). doi: 10.11144/Javeriana.ie20-1.epce
- Barros, L. M., Galindo Neto, N. M., Sá, G. G. M., Pereira, J. C. N., Barbosa, L. U., Oliveira Neto, J. G., ... Caetano, J. (2020). Experimental studies on COVID-19: overview of the worldscientific production. *Acta Paulista de Enfermagem*, 33, eAPE20200121. doi: 10.37689/actape/2020AO01215
- Bodys-Cupak, I., Majda, A., Skowron, J., Zaleska-Puchala, J., & Trzcinka, A. (2018). First year nursing students' coping strategies in stressful clinical practice situations. *Journal of Education in Science, Environment and Health*, 4(1), 12-18. Retrieved from <https://files.eric.ed.gov/fulltext/EJ1170988.pdf>
- Caldera-Villalobos, C., Garza-Veloz, I., Martínez-Avila, N., Delgado-Enciso, I., Ortiz-Castro, Y., Cabral-Pacheco, G. A., & Martínez-Fierro, M. L. (2020). The Coronavirus disease (COVID-19) challenge in Mexico: A critical and forced reflection as individuals and society. *Frontiers in Public Health*, 8, 337. doi: 10.3389/fpubh.2020.00337
- Cardona-Arias, J. A., Pérez-Restrepo, D., Rivera-Ocampo, S., Gómez-Martínez, J., & Reyes, Á. (2015). Prevalencia de ansiedad en estudiantes universitarios. *Diversitas: Perspectivas en Psicología*, 11(1), 79-89.
- Castillo, C., Chacón, T., & Díaz-Véliz, G. (2016). Ansiedad y fuentes de estrés académico en estudiantes de carreras de la salud. *Investigación en Educación Médica*, 5(20), 230-237. doi: 10.1016/j.riem.2016.03.001
- Chih-Cheng, L., Tzu-Ping, S., Wen-Chien, K., Hung-Jen, T., & Po-Ren, H. (2020). Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) and coronavirus disease-2019 (COVID-19): The epidemic and the challenges. *International Journal of Antimicrobial Agents*, 55(3), 105924. doi: 10.1016/j.ijantimicag.2020.105924

- Clemente, Z. A. (2021). "Estrés Académico y Ansiedad en Estudiantes de Licenciatura en Enfermería" [Tesis de Maestría] Benemérita Universidad Autónoma de Puebla, Facultad de Enfermería, Puebla, Mexico. Retrieved from <https://repositorioinstitucional.buap.mx/handle/20.500.12371/11807>
- Del Risco, M. C., Gómez, D. S., & Hurtado, F. S. (2015). "Estrés y rendimiento académico en estudiantes de la Facultad de enfermería-Universidad Nacional de la Amazonia Peruana, Iquitos-2015" [Tesis de Licenciatura]. Universidad Nacional de la Amazonia Peruana, Facultad de Enfermería, Perú. Retrieved from [https://repositorio.unapiquitos.edu.pe/bitstream/handle/20.500.12737/3954/Marie!%09Tesis\\_T%C3%ADtulo\\_2015.pdf.pdf?sequence=1&isAllowed=y](https://repositorio.unapiquitos.edu.pe/bitstream/handle/20.500.12737/3954/Marie!%09Tesis_T%C3%ADtulo_2015.pdf.pdf?sequence=1&isAllowed=y)
- Fandiño, M. I., Henríquez, N. M., & Rivera, D. E. (2020). Nivel de estrés y ansiedad en estudiantes de enfermería en práctica clínica de la Fundación Universitaria de San Gil Unisangil. Retrieved from <http://hdl.handle.net/20.500.12749/12423>
- Hernández, R. (2020). Estudio sobre la efectividad de un programa de intervención con refuerzo para reducir el estrés y la ansiedad en estudiantes de enfermería durante el Practicum [Tesis Doctoral]. Universidad Europea, Madrid, España. Retrieved from <https://www.educacion.gob.es/tesco/imprimirFicheroTesis.do?idFichero=sqU3bbJrkM4%3D>
- Lara, R. (2020). Educación en Enfermería en tiempos de pandemia: Desafíos presentes y futuros. *Ciencia y Enfermería (Impresa)*, 26. doi:10.29393/ce26-14eer110014
- López, J. M., Orbañanos, L., & Cibanal, M. L. (2013). Identificación de estresores en las prácticas clínicas y evolución de los mismos en una promoción de alumnos de enfermería. *Sanidad Militar*, 69(1). doi: 10.4321/S1887-85712013000100003
- Madeley, M. M. (2020). Estresores en las prácticas pre profesionales en internos de enfermería de Lima, 2020 [Tesis de Licenciatura]. Universidad César Vallejo, Lima, Perú. Retrieved from [https://repositorio.ucv.edu.pe/bitstream/handle/20.500.12692/58146/Marcos\\_MMG-SD.pdf?sequence=1&isAllowed=y](https://repositorio.ucv.edu.pe/bitstream/handle/20.500.12692/58146/Marcos_MMG-SD.pdf?sequence=1&isAllowed=y)
- Mahadeen, A., Abushaikha, L., & Habshneh, S. (2017). Educational experiences of undergraduate male nursing students: afocus group study. *Open Journal of Nursing*, 7, 50-57. doi: 10.4236/ojn.2017.71005
- Mamani, C. J., & Mamani, R. E. (2022). Estrés laboral y actitudes durante la pandemia COVID-19 del personal de enfermería a nivel de la micro-red de salud Cono Norte de Tacna, 2021. *Revista Muro de la Investigación*, 7(1). doi: 10.17162/rmi.v7i1.1690
- Moya, M., Larrosa, S., López, C., López, I., Morales, L., & Simón, Á. (2013). Percepción del estrés en los estudiantes de Enfermería ante sus prácticas clínicas. *Enfermería Global*, 12(31), 232-243. doi: 10.6018/eglobal.12.3.166261
- Rabelo-Silva, E. R., Monteiro, V., & Lumertz, M. A. (2022). Translación del conocimiento y avances en las prácticas de salud y enfermería. *Revista Gaúcha de Enfermagem*, 43(spe), e20220165. doi: 10.1590/1983-1447.2022.20220165.es
- Sánchez, A. M. (2019). Factores estresores en estudiantes de enfermería durante la realización de las practicas clínicas. *Nure Investigacion*, 16(99).
- Sánchez, F. T., & Talavera, M. F. (2021). Nivel de estrés y ansiedad en tiempos de COVID-19 en estudiantes del VIII y X semestre de la Facultad de Enfermería. Universidad Católica de Santa María, Arequipa 2020 [Tesis de Licenciatura]. Universidad Católica de Santa María, Arequipa, Perú. Retrieved from <https://www.nureinvestigacion.es/OJS/index.php/nure/article/view/1646>
- Tiga, D. C., Parra, D. I., & Domínguez, C. C. (2014). Competencias en proceso de enfermería en estudiantes de práctica clínica. *Revista Cuidarte*, 5(1), 585-594.
- Valero, N. J., Vélez, M. F., Duran, Á. A., & Torres, M. (2020). Afrontamiento del COVID-19: estrés, miedo, ansiedad y depresión. *Enfermería Investiga: Investigación, Vinculación, Docencia y Gestión*, 5(3), 63-70. doi: 10.31243/ei.uta.v5i3.913.2020
- Velásquez-Muñoz, A., Cornejo-Aravena, C., Bustamante-Toncio, F., & Mella-Guzmán, M. (2018). Estresores que influyen desfavorablemente en estudiantes chilenos de Obstetricia y Puericultura durante las practicas clínicas. *EDUMECENTRO*, 10(3), 12-24.
- Villanueva, L. E., & Ugarte, A. R. (2017). Niveles de ansiedad y la calidad de vida en estudiantes de una Universidad Privada de Arequipa. *Avances en Psicología: Revista de la Facultad de Psicología y Humanidades*, 25(2). doi: 10.33539/avpsicol.2017.v25n2.351